Study of Polarization Properties of Comet Halley Using a Mixture of Compact And Aggregate Particles

HIMADRI S. DAS¹

¹Department of Physics, Assam University, Silchar 788011, India. e-mail: hsdas@iucaa.ernet.in Phone: +91-94351-76013

The *in situ* measurements of comet Halley and the 'Stardust' returned samples of comet Wild 2 showed the presence of a mixture of silicates and organic refractory in cometary dust. In the present work, we propose a model which considers cometary dust as a mixture of aggregates and compact particles. We consider aggregates as ballistic cluster-cluster aggregate (BCCA) and ballistic agglomeration with one migration (BAM1) aggregate¹ and compact particles as spheroidal particles with some size distribution. For modeling comet 1P/ Halley, the power-law size distribution $n(a) \sim a^{-2.6}$, obtained from re-analysis of the Giotto spacecraft data², is taken. Using T-matrix code for polydisperse spheroids and Superposition T-matrix code for aggregates, the average simulated polarization curves are generated which can best fit to the observed polarization data at the three wavelengths $\lambda = 0.365 \mu m$, 0.485 μm and 0.684 μm . The best fitting complex refractive indices coming out from the present analysis correspond to a mixture of both silicates and organics, which are in good agreement with the *in situ* measurement of comet 1P/Halley by different spacecraft. The model successfully reproduces the positive part as well as the negative branch of the polarization at the above three wavelengths.

Keywords: Comet; dust; polarization; scattering; aggregates.

References

- [1] Y. Shen, B. T. Draine & E. T. Johnson, ApJ. 689, 260 (2008).
- [2] M. Fulle, A. C. Levasseur-Regourd, N. McBride, E. Hadamcik, 2000, AJ. 119, 1968 (2000).