

Measurement of Near-Infrared Diffuse Galactic Light

Kei Sano¹, Kimiaki Kawara¹, Shuji Matsuura², Hirokazu Kataza¹, Toshiaki Arai³,
Yoshiki Matsuoka⁴, Koji Tsumura³, Mai Shirahata³, Yosuke Onishi⁵

¹The University of Tokyo, Japan, ²Kwansei Gakuin University, Japan, ³Tohoku University, Japan,

⁴National Observatory of Japan, Japan, ⁵Tokyo Institute of Technology, Japan

We present new results of near-infrared diffuse Galactic light (DGL) from our recent study^{1,2,3}. The DGL consists of scattered light and thermal emission from interstellar dust grains illuminated by the interstellar radiation field. Therefore, the DGL measurement is useful in constraining some properties of interstellar dust, such as size distribution, albedo, and scattering asymmetry of dust grains.

In the diffuse interstellar medium in high Galactic latitudes, the DGL observation has been limited due to its faintness, particularly in the near-infrared wavelengths. We thus reanalyze all-sky maps obtained from Diffuse Infrared Background Experiment (DIRBE) onboard the *Cosmic Background Explorer* (COBE) satellite in the four near-infrared photometric bands (1.25, 2.2, 3.5, and 4.9 μm). As a result, we succeed in detecting the near-infrared DGL as a component that linearly correlates with interstellar 100 μm emission.

At 1.25 and 2.2 μm , our results are marginally consistent with the expected spectrum of scattered light assuming a recent interstellar dust model⁴. At 3.5 and 4.9 μm , thermal emission from stochastic heating of very small grains and fluorescence of polycyclic aromatic hydrocarbon (PAH) dominate the DGL. Compared with a recent thermal emission model in the diffuse interstellar medium⁵, we constrain the mass fraction of very small grains and PAH to the total dust to be more than $\sim 2\%$.

We also find that the intensity ratios of DGL to 100 μm emission are higher toward low Galactic latitudes at 1.25 and 2.2 μm . Since this trend is expected from the forward scattering characteristic of dust grains, we compare the obtained latitude dependence with the scattered light model taking into account the scattering asymmetry. The derived forward scattering characteristic is several times stronger than that expected from the recent dust model⁴. In addition to the scattered light component, latitude dependence of possible thermal emission in the diffuse interstellar medium may contribute to the obtained latitude dependence.

¹Sano et al. 2015, ApJ, 811, 77

²Sano et al. 2016, ApJ, 818, 72

³Sano et al. 2016, ApJL, 821, L11

⁴Weingartner & Draine 2001, ApJ, 548, 296

⁵Draine & Li 2007, ApJ, 657, 810